



Diabetic Retinopathy What You Should Know



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
National Institutes of Health
National Eye Institute

The National Eye Institute (NEI) conducts and supports research that leads to sight-saving treatments and plays a key role in reducing visual impairment and blindness. NEI is part of the National Institutes of Health, an agency of the U.S. Department of Health and Human Services.

For more information, contact—

National Eye Institute
National Institutes of Health
2020 Vision Place
Bethesda, MD 20892–3655
Telephone: 301–496–5248
Email: 2020@nei.nih.gov
Website: www.nei.nih.gov



About Diabetic Retinopathy

1



Detection

3



Prevention and Treatment

5



What You Can Do

10



Current Research

12



Additional Resources


13





What is Diabetic Retinopathy?

Diabetic retinopathy is a complication of diabetes and the leading cause of vision impairment and blindness among working-age adults. It occurs when diabetes damages the tiny blood vessels in the retina, which is the light-sensitive tissue at the back of the eye. Diabetic retinopathy may lead to diabetic macular edema (DME), which is a swelling in an area of the retina called the macula.



Diabetic retinopathy involves damage to the retina, the light-sensitive tissue at the back of the eye.

What causes diabetic retinopathy?

Chronically high blood sugar from diabetes is associated with damage to the tiny blood vessels in the retina, leading to diabetic retinopathy. The retina detects light and converts it to signals sent through the optic nerve to the brain. Diabetic retinopathy can cause blood vessels in the retina to leak fluid

or hemorrhage (bleed), distorting vision. In its most advanced stage, new abnormal blood vessels proliferate (increase in number) on the surface of the retina, which can lead to scarring and cell loss in the retina.

Diabetic retinopathy may progress through four stages:

- 1. Mild nonproliferative retinopathy.** Small areas of balloon-like swelling in the retina's tiny blood vessels, called microaneurysms, occur at this earliest stage of the disease. These microaneurysms may leak fluid into the retina.
- 2. Moderate nonproliferative retinopathy.** As the disease progresses, blood vessels that nourish the retina may swell and distort. They may also lose their ability to transport blood. Both conditions cause characteristic changes to the appearance of the retina and may contribute to DME.
- 3. Severe nonproliferative retinopathy.** Many more blood vessels are blocked, depriving blood supply to areas of the retina. These areas secrete growth factors that signal the retina to grow new blood vessels.
- 4. Proliferative diabetic retinopathy (PDR).** At this advanced stage, growth factors secreted by the retina trigger the proliferation of new blood vessels, which grow along the inside surface of the retina and into the vitreous gel, the fluid that fills the eye. The new blood vessels are fragile, which makes them more likely to leak and bleed. Accompanying scar tissue can contract and cause retinal detachment—the pulling away of the retina from underlying tissue, like wallpaper peeling away from a wall. Retinal detachment can lead to permanent vision loss.



What is diabetic macular edema (DME)?

DME is the build-up of fluid (edema) in a region of the retina called the macula. The macula is important for the sharp, straight-ahead vision that is used for reading, recognizing faces, and driving. DME is the most common cause of vision loss among people with diabetic retinopathy. About half of all people with diabetic retinopathy will develop DME. Although it is more likely to occur as diabetic retinopathy worsens, DME can happen at any stage of the disease.

Who is at risk for diabetic retinopathy?

People with all types of diabetes (type 1, type 2, and gestational) are at risk for diabetic retinopathy. Risk increases the longer a person has diabetes. Between 40 and 45 percent of Americans diagnosed with diabetes have some stage of diabetic retinopathy, although only about half are aware of it. Women who develop or have diabetes during pregnancy may have rapid onset or worsening of diabetic retinopathy.

Detection

What are the symptoms of diabetic retinopathy and DME?

The early stages of diabetic retinopathy usually have no symptoms. The disease often progresses unnoticed until it affects vision. Bleeding from abnormal retinal blood vessels can cause the appearance of "floating" spots. These spots sometimes clear on their own. But without prompt treatment, bleeding often recurs, increasing the risk of permanent vision loss. If DME occurs, it can cause blurred vision.



The same scene as viewed by a person normal vision (left) and with (center) advanced diabetic retinopathy. The floating spots are hemorrhages that require prompt treatment. DME (right) causes blurred vision.

How are diabetic retinopathy and DME detected?

Diabetic retinopathy and DME are detected during a comprehensive dilated eye exam that includes:

- 1. Visual acuity testing.** This eye chart test measures a person's ability to see at various distances.
- 2. Tonometry.** This test measures pressure inside the eye.
- 3. Pupil dilation.** Drops placed on the eye's surface dilate (widen) the pupil, allowing a physician to examine the retina and optic nerve.
- 4. Optical coherence tomography (OCT).** This technique is similar to ultrasound but uses light waves instead of sound waves to capture images of tissues inside the body. OCT provides detailed images of tissues that can be penetrated by light, such as the eye.

A comprehensive dilated eye exam allows the doctor to check the retina for:



1. Changes to blood vessels
2. Leaking blood vessels or warning signs of leaky blood vessels, such as fatty deposits
3. Swelling of the macula (DME)
4. Changes in the lens
5. Damage to nerve tissue

If DME or severe diabetic retinopathy is suspected, a **fluorescein angiogram** may be used to look for damaged or leaky blood vessels. In this test, a fluorescent dye is injected into the bloodstream, often into an arm vein. Pictures of the retinal blood vessels are taken as the dye reaches the eye.

Prevention and Treatment

How can people with diabetes protect their vision?

Vision lost to diabetic retinopathy is sometimes irreversible. However, early detection and treatment can reduce the risk of blindness by 95 percent. Because diabetic retinopathy often lacks early symptoms, people with diabetes should get a comprehensive dilated eye exam at least once a year. People with diabetic retinopathy may need eye exams more frequently. Women with diabetes who become pregnant should have a comprehensive dilated eye exam as soon as possible. Additional exams during pregnancy may be needed.

Studies such as the Diabetes Control and Complications Trial (DCCT) have shown that controlling diabetes slows the onset and worsening of diabetic retinopathy. DCCT study participants who kept their blood glucose level as close to normal as possible were significantly less likely than those without optimal glucose control to develop diabetic retinopathy, as well as kidney and nerve diseases. Other trials have shown that controlling elevated blood pressure and cholesterol can reduce the risk of vision loss among people with diabetes.

Treatment for diabetic retinopathy is often delayed until it starts to progress to PDR, or when DME occurs. Comprehensive dilated eye exams are needed more frequently as diabetic retinopathy becomes more severe. People with severe nonproliferative diabetic retinopathy have a high risk of developing PDR and may need a comprehensive dilated eye exam as often as every 2 to 4 months.

How is DME treated?

DME can be treated with several therapies that may be used alone or in combination.


Anti-VEGF Injection Therapy. Anti-VEGF drugs are injected into the vitreous gel to block a protein called vascular endothelial growth factor (VEGF), which can stimulate abnormal blood vessels to grow and leak fluid. Blocking VEGF can reverse abnormal blood vessel growth and decrease fluid in the retina. Available anti-VEGF drugs include Avastin (bevacizumab), Lucentis (ranibizumab), and Eylea (aflibercept). Lucentis and Eylea are approved by the U.S. Food and Drug Administration (FDA) for treating DME.



Avastin was approved by the FDA to treat cancer, but is commonly used to treat eye conditions, including DME.

The NEI-sponsored Diabetic Retinopathy Clinical Research Network compared Avastin, Lucentis, and Eylea in a clinical trial. The study found all three drugs to be safe and effective for treating most people with DME. Patients who started the trial with 20/40 or better vision experienced similar improvements in vision no matter which of the three drugs they were given. However, patients who started the trial with 20/50 or worse vision had greater improvements in vision with Eylea.

Most people require monthly anti-VEGF injections for the first six months of treatment. Thereafter, injections are needed less often: typically three to four during the second



The retina of a person with diabetic retinopathy and DME, before (left) and after anti-VEGF treatment (right). The large white circle is the optic nerve. The bright yellow spots are fatty deposits. Note that they have mostly cleared after treatment.

six months of treatment, about four during the second year of treatment, two in the third year, one in the fourth year, and none in the fifth year. Dilated eye exams may be needed less often as the disease stabilizes.

Avastin, Lucentis, and Eylea vary in cost and in how often they need to be injected, so patients may wish to discuss these issues with an ophthalmologist.

Focal/grid macular laser surgery. In focal/grid macular laser surgery, a few to hundreds of small laser burns are made to leaking blood vessels in areas of edema near the center of the macula. Laser burns for DME slow the leakage of fluid, reducing swelling in the retina. The procedure is usually completed in one session, but some people may need more than one treatment. Focal/grid laser is sometimes applied before anti-VEGF injections, sometimes on the same day or a few days after an anti-VEGF injection, and sometimes only when DME fails to improve adequately after six months of anti-VEGF therapy.

Corticosteroids. Corticosteroids, either injected or implanted into the eye, may be used alone or in combination with other drugs or laser surgery to treat DME. The Ozurdex (dexamethasone) implant is for short-term use, while the Iluvien (fluocinolone acetonide) implant is longer lasting. Both are biodegradable and release a sustained dose of corticosteroids to suppress DME. Corticosteroid use in the eye increases the risk of cataract and glaucoma. DME patients who use corticosteroids should be monitored for increased pressure in the eye and glaucoma.



How is proliferative diabetic retinopathy (PDR) treated?

For decades, PDR has been treated with scatter laser surgery, sometimes called panretinal laser surgery or panretinal photocoagulation. Treatment involves making 1,000 to 2,000 tiny laser burns in areas of the retina away from the macula. These laser burns are intended to cause abnormal blood vessels to shrink. Although treatment can be completed in one session, two or more sessions are sometimes required. While it can preserve central vision, scatter laser surgery may cause some loss of side (peripheral), color, and night vision. Scatter laser surgery works best before new, fragile blood vessels have started to bleed. Recent studies have shown that anti-VEGF treatment not only is effective for treating DME, but is also effective for slowing progression of diabetic retinopathy, including PDR, so anti-VEGF is increasingly used as a first-line treatment for PDR.

What is a vitrectomy?

A vitrectomy is the surgical removal of the vitreous gel in the center of the eye. The procedure is used to treat severe bleeding into the vitreous, and is performed under local or general anesthesia. Ports (temporary water-tight openings) are placed in the eye to allow the surgeon to insert and remove instruments, such as a tiny light or a small vacuum called a vitrector. A clear salt solution is gently pumped into the eye through one of the ports to maintain eye pressure

during surgery and to replace the removed vitreous. The same instruments used during vitrectomy also may be used to remove scar tissue or to repair a detached retina.

Vitrectomy may be performed as an outpatient procedure or as an inpatient procedure, usually requiring a single overnight stay in the hospital. After treatment, the eye may be covered with a patch for days to weeks and may be red and sore. Drops may be applied to the eye to reduce inflammation and the risk of infection. If both eyes require vitrectomy, the second eye usually will be treated after the first eye has recovered.

What You Can Do

What should I ask my eye care professional?

You can protect yourself against vision loss by working in partnership with your eye care professional. Ask questions and get the information you need to take care of yourself and your family.

What are some questions to ask?

About my eye disease or disorder...

- What is my diagnosis?
- What caused my condition?
- Can my condition be treated?
- How will this condition affect my vision now and in the future?



- Should I watch for any particular symptoms and notify you if they occur?
- Should I make any lifestyle changes?

About my treatment...

- What is the treatment for my condition?
- When will the treatment start and how long will it last?
- What are the benefits of this treatment and how successful is it?
- What are the risks and side effects associated with this treatment?
- Are there foods, drugs, or activities I should avoid while I'm on this treatment?
- If my treatment includes taking medicine, what should I do if I miss a dose?
- Are other treatments available?

About my tests...

- What kinds of tests will I have?
- What can I expect to find out from these tests?
- When will I know the results?
- Do I have to do anything special to prepare for any of the tests?
- Do these tests have any side effects or risks?
- Will I need more tests later?

What if treatment doesn't improve vision?

An eye care professional can help locate and make referrals to low vision and rehabilitation services and suggest devices that may help make the most of remaining vision. Many community organizations and agencies offer information about low vision counseling, training, and other special services for people with visual impairment. A nearby school of medicine or optometry also may provide low vision and rehabilitation services.

What research is being done?

The NEI is conducting and supporting research that seeks better ways to detect, treat, and prevent vision loss in people with diabetes. This research is being conducted in labs and clinical centers across the country.

For example, the Diabetic Retinopathy Clinical Research Network (DRCR.net) conducts large multi-center trials to test new therapies for diabetic eye disease, and to compare different therapies. The network formed in 2002 and comprises more than 350 physicians practicing at more than 140 clinical sites across the country. Many of the sites are private practice eye clinics, enabling the network to quickly bring innovative treatments from research into community practice.

NEI-funded scientists are also seeking ways to detect diabetic retinopathy at earlier stages. For example, researchers are harnessing a technology called adaptive



optics (AO) to improve imaging techniques such as OCT. AO was developed by astronomers seeking to improve the resolution of their telescopes by filtering out distortions in the atmosphere. In the clinic, diagnostic devices that use AO may improve the detection of subtle changes in retinal tissue and blood vessels.

Where can I get more information?

For more information about diabetic retinopathy, you may wish to contact:

American Academy of Ophthalmology

P.O. Box 7424

San Francisco, CA 94120-7424

415-561-8500

www.aao.org

www.eyecareamerica.org (Online Referral)

American Diabetes Association

1701 North Beauregard Street

Alexandria, VA 22311

1-800-342-2383 (National Headquarters)

703-549-1500

1-888-342-2383 (Washington DC Office)

Email: askada@diabetes.org

www.diabetes.org

American Foundation for the Blind

2 Penn Plaza, Suite 1102

New York, NY 10121

1-800-232-5463

212-502-7600

E-mail: afbinfo@afb.net

www.afb.org

American Optometric Association

243 North Lindbergh Boulevard, Floor 1

St. Louis, MO 63141-7851

314-991-4100

Toll-free (800) 365-2219

www.aoa.org

Council of Citizens with Low Vision International

2200 Wilson Blvd. Suite 650

Arlington, VA 22201

1-800-733-2258

Email: president@cclvi.org

www.cclvi.org

Juvenile Diabetes Research Foundation International

26 Broadway

New York, NY 10004

1-800-553-CURE (2873)

(212) 785-9500

Fax: (212) 785-9595

Email: info@jdrf.org

www.jdrf.org



National Eye Institute

National Institutes of Health
2020 Vision Place
Bethesda, MD 20892-3655
301-496-5248
Email: 2020@nei.nih.gov
www.nei.nih.gov

National Institute of Diabetes and Digestive and Kidney Diseases Clearinghouses Publications Catalog

5 Information Way
Bethesda, MD 20892-3568
1-800-860-8747
TTY: 1-866-569-1162
Fax: 301-634-0716
Email: catalog@niddk.nih.gov
www.niddk.nih.gov

Prevent Blindness

211 West Wacker Drive, Suite 1700
Chicago, IL 60606
1-800-331-2020
Email: info@preventblindness.org
www.preventblindness.org

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
National Institutes of Health
National Eye Institute
Revised 9/15



National Eye Institute